

Know Your Steps:

The A, B, C's of Successful Injection

epinephrine injection, USP
auto-injector

Available as

0.15 mg

0.3 mg

STEP A

- Pull off **GRAY** end cap labeled [1]; you will now see a **RED** tip. Never put thumb, finger, or hand over the **RED** tip.
- Pull off **GRAY** end cap with [2].



STEP B

- Put the **RED** tip against the middle of the outer side of your thigh as shown.
- Press down hard and hold in place while slowly counting to 10.



STEP C

- Call **911**. Seek immediate medical attention.



Indication

- Epinephrine auto-injector is an emergency injection ("shot") of epinephrine. It is a medicine used for life-threatening allergic reactions such as severe swelling, breathing problems, or loss of blood pressure. Allergic reactions can be caused by stinging and biting insects (bugs), allergy injections, food, medicines, exercise, or unknown causes.
- After using epinephrine auto-injector go to your doctor or emergency room right away for more medical treatment.

Important Safety Information (Also see reverse side)

- Too much epinephrine (epinephrine auto-injector) can cause dangerously high blood pressure, stroke or death.
- If you take certain medicines, you may develop serious life-threatening side effects from the epinephrine in epinephrine auto-injector. Make sure to tell your doctor about all your medical conditions and allergies.

Epinephrine can cause the following reactions:

- ♦ faster, irregular (wrong) or "pounding" heartbeat
- ♦ throbbing headache
- ♦ paleness
- ♦ feelings of over excitement, anxiety, or fear
- ♦ weakness or shakiness
- ♦ dizziness
- ♦ nausea and vomiting
- ♦ sweating

- Some reactions can be serious. They usually go away with rest.
- These are not all the possible side effects of epinephrine auto-injector. For more information, ask your doctor or pharmacist.

You are encouraged to report side effects of prescription drugs to the FDA by visiting www.FDA.gov/medwatch or calling 1-800-FDA-1088. Ask your healthcare provider for additional information about Lineage Therapeutics EPINEPHRINE AUTO-INJECTOR and anaphylaxis.

Epinephrine injection, USP auto-injector is an authorized generic of Adrenaclick® (epinephrine injection, USP), a trademark of Amedra Pharmaceuticals LLC.

Please see package insert for full prescribing information, available at www.epinephrineautoinject.com.

To learn more about epinephrine auto-injector go to
www.epinephrineautoinject.com

 lineage therapeutics

JUNE2013029. © 2013.
Lineage Therapeutics.
All rights reserved.

Indication

- Epinephrine auto-injector is an emergency injection (“shot”) of epinephrine. It is a medicine used for life-threatening allergic reactions such as severe swelling, breathing problems, or loss of blood pressure. Allergic reactions can be caused by stinging and biting insects (bugs), allergy injections, food, medicines, exercise, or unknown causes.
- After using epinephrine auto-injector go to your doctor or emergency room right away for more medical treatment.

Important Safety Information

- Make sure to tell your doctor about all your medical conditions and allergies, but especially if you:
 - ◆ have heart disease or high blood pressure
 - ◆ have diabetes
 - ◆ have thyroid conditions
 - ◆ have asthma
 - ◆ have depression or other mental disease
 - ◆ have Parkinson’s disease
 - ◆ are pregnant
 - ◆ are allergic to any of the ingredients in epinephrine auto-injector
- Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins and herbal supplements. Some medicines may cause serious side effects if taken while you use epinephrine auto-injector. Some medicines may affect how epinephrine auto-injector works, or epinephrine auto-injector may affect how your other medicines work.
- Diabetic patients may need to adjust the dose of their diabetes medicines or insulin after using epinephrine auto-injector.
- Avoid injecting epinephrine auto-injector into your buttock or any other part of your body, other than the middle of the outer side of your thigh (upper leg). Avoid injecting epinephrine auto-injector into a vein.
- Too much epinephrine can cause dangerously high blood pressure, stroke, or death. If you take certain medicines, you may develop serious life-threatening side effects from the epinephrine

in epinephrine auto-injector. Be sure to tell your doctor about all the medicines you take, especially medicines for asthma.

- Patients with certain medical conditions, or who take certain medicines, may get more side effects from epinephrine auto-injector, or the side effects may last longer. This includes patients who take certain types of medicines for asthma, allergies, depression, hyperthyroidism, high blood pressure, and heart disease. Patients with heart disease may feel chest pain (angina). Patients with mental disease or Parkinson’s disease may have worsening symptoms of their illness. Epinephrine can cause the following reactions:
 - ◆ faster, irregular (wrong) or “pounding” heartbeat
 - ◆ throbbing headache
 - ◆ paleness
 - ◆ feelings of over excitement, anxiety, or fear
 - ◆ weakness or shakiness
 - ◆ dizziness
 - ◆ nausea and vomiting
 - ◆ sweating
- Some reactions can be serious. They usually go away with rest.
- These are not all the possible side effects of epinephrine auto-injector. For more information, ask your doctor or pharmacist.
- You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

AFTER USE/DISPOSAL

- The remaining liquid that is left after this fixed dose cannot be further administered and should be discarded.
- Put the auto-injector, needle-first, into the carrying case.
- Put the light gray case top over the non-needle end of the epinephrine auto-injector unit and close the case.
- Give your used epinephrine auto-injector to a healthcare worker for proper disposal. Do NOT throw away in a regular trash can.